



Head and Neck Associates of Orange County, Inc.

An Incorporated Medical Group

Head & Neck Surgery

Pediatric & Adult Otolaryngology

Facial Reconstructive Surgery

NASAL AIRWAY SURGERY SURGICAL INSTRUCTIONS

INTRODUCTION

Your doctor has recommended nasal surgery for you. The following information is provided to help you prepare for your surgery, and to help you understand more clearly the associated benefits, risks, and complications of nasal surgery. You are encouraged to ask your doctor any questions that you feel necessary to help you better understand the above procedure.

Nasal obstruction is usually caused by a deviated or crooked septum or enlarged tissues (turbinates) within the nose. A septoplasty is an operation to correct a deformity of the partition (the septum) between the two sides of the nose. A turbinectomy is the surgical removal of an abnormally enlarged turbinate from inside the nose to improve breathing. A turbinectomy can be done by removal of tissue (sub-mucus resection) or by radio-frequency (coblation). The goals of nasal airway surgery include the following: To decrease nasal obstruction and improve airflow through the nose, to control nosebleeds, to enhance visualization of the inside of the nose, to relieve nasal headaches associated with swelling of the inside of the nose, and to promote drainage of the sinus cavities. Nasal airway surgery is done through the nostrils without the need for external incisions.

The following instructions are designed to help you recover from nasal surgery as easily as possible. Taking care of you can prevent complications. It is very important that you read these instructions and follow them carefully. We will be happy to answer any questions.

RISKS AND COMPLICATIONS

Your surgery will be performed safely and with care in order to obtain the best possible results. You have the right to be informed that the surgery may involve risks of unsuccessful results, complications, or injury from both known and unforeseen causes. Because individuals differ in their response to surgery, their anesthetic reactions, and their healing outcomes, ultimately there can be no guarantee made as to the results or potential complications. Furthermore, surgical outcomes may be dependent on preexisting or concurrent medical conditions.

The following complications have been reported in the medical literature. This list is not meant to be inclusive of every possible complication. They are listed here for your information only, not to frighten you, but to make you aware and more knowledgeable concerning this surgical procedure. Although many of these complications are rare, all have occurred at one time or another in the hands of experienced surgeons practicing the standards of community care. Anyone who is contemplating surgery must weigh the potential risks and complications against the potential benefits of the surgery, or any alternative to surgery.

Initials _____

1. Nasal obstruction due to the failure to straighten the septum or later re-deviation of the septum, collapse of the nasal cartilage's, or the re-growth or swelling of the turbinates.
2. Failure to resolve coexisting sinus infections, or recurrence of coexisting sinus problems and/or polyps, or the need for further or more aggressive surgery.
3. Bleeding. In rare situations, a need for blood products or a blood transfusion. You have the right, should you choose, to have autologous or designated donor directed blood prepared in advance in case an emergency transfusion was necessary. You are encouraged to consult with your doctor if you are interested in this option.
4. Chronic nasal drainage or excessive dryness or crusting of the nose or sinuses.
5. Need for allergy evaluation, treatments, or environmental controls. Surgery is neither a cure for nor a substitute for good allergy control or treatment.
6. Failure to improve or resolve concurrent respiratory illness such as, but not limited to asthma, bronchitis, or cough.
7. Failure to resolve associated "sinus or nasal" headaches. The exact cause of headaches can be difficult to determine or be multi-factorial. You may require consultation with another specialist such as a neurologist.
8. Permanent numbness of the upper teeth.
9. Prolonged pain, impaired healing, and the need for hospitalization.
10. Septal perforation, which is a permanent hole inside the nose between the two sides. In rare situations this may change the outer appearance of the nose.
11. Failure to restore or worsening of the sense of smell or taste, or failure to relieve nosebleeds.

BEFORE SURGERY

In most situations, the surgery is performed at the hospital or at the outpatient surgery center. An anesthesiologist will monitor you throughout the procedure. Usually, the anesthesiologist will call the night before surgery to review the medical history. If he or she is unable to reach you the night before surgery, they will talk with you that morning. If your doctor has ordered preoperative laboratory studies, you should arrange to have these done several days in advance. Arrange for someone to pick you up from the surgical facility and to spend the first night after surgery with you.

You should not take aspirin, or any product containing aspirin, within 10 days of the date of your surgery. Non-steroidal anti-inflammatory medications (such as Advil) should not be taken within 7 days of the date of surgery. Many over-the-counter products contain aspirin or Advil type drugs so it is important to check all medications carefully. If there is any question please call the office or consult your Pharmacist. Tylenol is an acceptable pain reliever. Usually your doctor will give you your prescriptions at the preoperative visit. It is best to have these filled prior to the date of your surgery. Smokers should make every effort to stop smoking, or at least reduce the number of cigarettes. This will help to reduce postoperative coughing and bleeding, and to improve healing.

You must not eat or drink anything 6 hours prior to the time of surgery. This includes even water, candy, or chewing gum. Anything in the stomach increases the chances of an anesthetic complication.

Nasal Airway

Initials _____

Smokers should make every effort to stop smoking, or at least reduce the number of cigarettes. This will help to improve the healing process and to reduce postoperative coughing and bleeding. **It is important that you stop using Afrin or other OTC(over the counter) nasal sprays at least 1 week prior to surgery. (Saline sprays are OK) If you cannot breathe through your nose without using these OTC sprays notify your doctor BEFORE surgery.**

If you are sick or have a fever the day before surgery, call the office. If you wake up sick the day of surgery, still proceed to the surgical facility as planned. Your doctor will decide if it's safe to proceed with surgery.

After surgery, your doctor will usually like for you to start saline irrigation of the nose and sinuses. It is best to use a water pick for these irrigation's. You will need to either purchase or borrow a water pick. Your doctor will instruct you on how and when to start using your water pick irrigator. He will also give you a special device to attach to the water pick to make it easier to use (there will be a \$20 extra charge for this attachment).

THE DAY OF SURGERY

It is important that you know precisely what time you are to check in with the surgical facility and that you allow sufficient preparation time. Bring all papers and forms with you including any preoperative orders and history sheets.

You should wear comfortable loose fitting clothes that do not have to be pulled over your head. Leave all jewelry and valuables at home. Remove all make-up with a cleansing cream. Thoroughly wash your face and neck with soap and water. Do not apply make-up or cream to your face.

Do not take any medication unless instructed by your doctor or the anesthesiologist. Usually in the pre-operative holding room, a nurse will start an intravenous infusion line (IV) and you may be given a medication to help you relax.

DURING SURGERY

In the operating room, the anesthesiologist will usually use a mixture of a gas and an intravenous medication to put you to sleep and to maintain your anesthetic at a safe and comfortable level. During the procedure, you will be continuously monitored including pulse oximeter (oxygen saturation) and cardiac rhythm (EKG). The surgical team is prepared for any emergency. In addition to the surgeons and the anesthesiologist, there will be a nurse and a surgical technician in the room.

The whole procedure usually takes about one hour. Your doctor will come to the waiting room to talk with any family or friends once you are safely to the recovery room.

Nasal Airway

Initials_____

AFTER SURGERY

After surgery, you will be taken to the recovery room where a nurse will monitor you for about one hour. You will need to arrange to have a friend or family member to pick you up from the surgical facility and to take you home. He or she should spend the first night with you after your return home.

When you go home you should go to bed and rest with your head elevated on 2-3 pillows. By keeping your head elevated above your heart, you can minimize edema and swelling. You may get out of bed with assistance to use the bathroom. Avoid straining, if you are constipated, take a stool softener or a gentle laxative.

It is best to eat a light, soft, and cool diet as tolerated once you have recovered fully from the anesthetic. Even though you may be hungry immediately after surgery, it is best to go slowly to prevent postoperative nausea and vomiting. Occasionally, you may vomit one or two times immediately after surgery; if it persists, your doctor may prescribe medications to settle the stomach. It is important to remember that a good overall diet with ample rest promotes healing.

You may have some swelling and bruising of your nose, upper lip, cheeks, or around your eyes for several days after surgery. This swelling is normal, and will gradually go away over the next 7 to 10 days. You can help reduce it by keeping ice on your face, bridge of the nose, and eyes as much as tolerated. This will also help with postoperative edema and pain. Some patients have found frozen vegetable in packages (for example bags of frozen peas) to be a convenient ice pack which is more likely to conform to the face.

Moderate bleeding from the nose is normal, and will gradually decrease. The gauze dressing ("mustache dressing") will collect blood and should be changed only when saturated. It is not unusual to change the dressing every hour during the first 24 hours after surgery. After a few days you will probably no longer need to use the dressing. Do not take aspirin, aspirin-containing medications, or non-steroidal anti-inflammatory medications (Advil) for 3 weeks following surgery.

You will likely be prescribed antibiotics after surgery, and should finish all the pills that have been ordered. Some form of a narcotic will also be prescribed (usually Vicodin), and is to be taken as needed. If you require narcotics you are cautioned not to drive. If you have nausea or vomiting postoperatively, you may be prescribed anti-emesis medications such as Phenergan. If you have any questions or you feel that you are developing a reaction to any of these medications, you should consult your doctor. You should not take any other medication, either prescribed or over-the-counter, unless you have discussed it with your doctor.

GENERAL INSTRUCTIONS AND FOLLOW-UP CARE

In most situations, packs will have been placed in your nose to control postoperative bleeding. Your doctor will tell you when you are expected to return to the office to have these removed.

Nasal Airway

Initials_____

You need to call the office to schedule this postoperative appointment. Please arrange for someone to drive you to and from the office for this first visit. You should eat a light meal before coming, and avoid taking excessive pain medications. You will also have several subsequent office visits to assess healing, remove crusts, and insure a speedy recovery. Your nose will probably be tender after surgery, so we will spray your nose with a special numbing medication before removing crusts. Immediately after the nasal packs are removed you should use a saline nasal spray such as "Ocean Spray" several times per day to prevent crusts from forming in your nose.

After the packing has been removed, you may breathe through your nose, but do not blow or sneeze through it for 7-10 days. If you must sneeze, open your mouth. Expect some light blood-tinged drainage from the nose for several days. If bleeding becomes excessive, apply ice and rest quietly with the head elevated while holding your nose. If bleeding continues, call the office.

One of the most important things you can do after surgery is nasal irrigation. Immediately after the nasal packs are removed you should use a saline nasal spray such as "Ocean Spray" several times per day to prevent crusts from forming in your nose. We will also have you start using a water pick irrigator or NeilMed Sinus Rinse Kit during the first week of surgery, depending on your surgeon. You are to continue using the nasal irrigator until we tell you to stop. Many patients will use it indefinitely or when they have an upper respiratory infection. Failure to irrigate will likely cause the surgery to fail!!

You may go back to work or school only when your doctor gives you medical clearance. You are encouraged to rest for the first week following surgery. Avoid excessive talking, smiling, hard chewing, strenuous activities, lifting heavy objects, and bending over. Alcohol and tobacco should be avoided because they may prolong swelling and healing.

Use a Q-tip to lubricate your nostrils with Vaseline; hydrogen peroxide with help to loosen crusts. After several days you may notice a few of the absorbable sutures. Be gentle while brushing your upper teeth. They will often be tender for several weeks, and you may have some numbness of the teeth and palate for several months. You may use make-up at any time to camouflage any bruising.

After 3 weeks, if you are not having problems with bleeding, you may resume exercise and swimming, but no diving for two months. You should plan to stay in town for 3 weeks to allow for postoperative care.

NOTIFY YOUR DOCTOR IF YOU HAVE

1. A sudden increase in the amount of bruising and pain associated with excessive swelling of the nose and possible difficulty breathing.
2. A fever greater than 101.5 degrees which is persistent despite increasing the amount of fluid you drink and Tylenol. A person with a fever should try to drink approximately one cup of fluid each waking hour.

Nasal Airway

Initials_____

3. Persistent sharp pain or headache which is not relieved by the prescribed pain medication.
4. Increased swelling or redness of the nose or eyes.

IF YOU WOULD LIKE TO LEARN MORE

The physicians of Head and Neck Associates recommend **www.medicinenet.com** as an internet source of information. MedicineNet® is a network of U.S. Board Certified Physicians and Allied Health Professionals. Find easy-to-understand medical information to make smart health decisions with your doctor. Stay informed on all aspects of health and medicine. Get the latest health and medical news delivered to your e-mail box!

Please visit our web site at **www.hnaoc.com**

SURGICAL FINANCIAL POLICY

Head and Neck Associates of Orange County will submit claims to your insurance company for any surgical procedures performed by our physicians. Prior to your scheduled surgery date our staff will verify eligibility and benefits. We will also obtain precertification and/ or authorization when required by your insurance company. Please be aware however, that this is not a guaranty of payment. Any expenses deemed not covered by your insurance company for any reason, will be your financial responsibility. All monies owed by the patient, i.e., remaining deductible or coinsurance amounts and any procedures or fees deemed not medically necessary, are due prior to the date of surgery. All financial arrangements must be made prior to the date of surgery. Please be aware that this office will bill only for the physicians' services. Any other services related to your surgery, i.e., facility, anesthesiology, radiology, laboratory or pathology will be billed by the facility providing these services, and not included in our billings.

Our office accepts the following forms of payment; Visa or Mastercard, cash and personal checks. A twenty dollar service charge will be assessed to your account for any check returned by your bank.

IMPORTANT PHONE NUMBERS

If you have any questions, do not hesitate to call the office at (949) 364-4361. At night or on the weekends, if your need is urgent and cannot wait until regular business hours, you may call our emergency number at (949) 470-1403. Our answering service will contact the doctor on call. If for some reason you cannot reach the doctor on call and your need is life threatening, go to the emergency room.

Nasal Airway

Initials _____

